

Categories of Medications Specifically of Concern When Performing Myelograms

MAO INHIBITORS (Hold 48 hours prior to exam, resume next day)	SSRI's (SELECTIVE SEROTONIN REUPTAKE INHIBITORS) (Hold 48 hours prior to exam, resume next day)
Deprenyl (selegiline) Eldepryl (selegiline) Marplan (isocarboxazid) Nardil (phenelzine) Parnate (tranylcypramine)	Celexa (citalopram) Cymbalta (duloxetine) Lexapro (escitalopram oxalate) (Lexapril) Luvox (fluvoxamine) Paxil (paroxetine) Prozac (fluoxetine) (Sarafem) Zoloft (sertraline)
OTHER ANTIDEPRESSANTS (Hold 48 hours prior to exam, resume next day)	TRICYCLIC ANTIDEPRESSANTS (Hold 48 hours prior to exam, resume next day)
Buspar (buspirone) Cymbalta (duloxetine) Desyrel (trazodone) Edronax, Vestra (reboxetine) Effexor (venlafaxine) Ludiomil (maprotiline) Remeron (mirtazapine) Serzone (nefazodone) Wellbutrin (bupropion) Desvenlafaxine (Pristiq) Milnacipram (Savella)	Adapin (doxepin) Anafranil (clomipramine) Asendin (amoxapine) Elavil (amitriptyline) Endep (amitriptyline) Entrafon (amitriptyline and perphenazine) Janimine (imipramine) Norpramin (desipramine) Pamelor (Nortriptyline) Pertofrane (desipramine) Sinequan (doxepin) Surmontil (trimipramine) Tofranil (imipramine) Triavil (amitriptyline and perphenazine) Trilafon (amitriptyline and perphenazine) Vivactil (protriptyline)

Medications of concern when Performing Myelograms Dec.2004

Revised Feb. 2005, June 2005, October 2005, November 2005, June 2006, Oct 2006, July 2009 (DKS)

<p style="text-align: center;">PHENOTHIAZINES (Hold 48 hours prior to exam, resume next day)</p>	<p style="text-align: center;">ANTIPSYCHOTICS (Hold 48 hours prior to exam, resume next day)</p>
<p>Compazine (prochlorperazine) Entrafon (amitriptyline and perphenazine) Flexeril (cyclobenzaprine) Mellaril (thioridazine) Mepergan (promethazine and meperidine) (Phenothiazine and Narcotic) Phenergan (Promethazine) Prolixin (fluphenazine) Prozine (promethazine) Serentil (mesoridazine) Sparine (promethazine) Stelazine (trifluoperazine) Synalgos-DC (promethazine and aspirin and caffeine) (Phenothiazine and Anti-inflammatory and CNS Stimulant) Tacaryl (methdilazine) Temaril (trimeprazine) Thorazine (chlorpromazine) Triavil (amitriptyline and perphenazine) Trilafon (amitriptyline and perphenazine)</p>	<p>Carbolith (lithium) Clozaril (clozapine) Haldol (haloperidol) Inapsine (droperidol) Innovar (droperidol and fentanyl) (Anti-psychotic and Narcotic) Lithane (lithium) Loxitane (loxapine) Moban (molindone) Navane (thiothixene) Olanzapine (zyprexa) (zydis) Orap (pimozide) Seroquel(quetiapine fumarate) Risperidone (Risperdal) Iloperidone (Fanapt) Paliperidone (Invega)</p> <p style="text-align: center;">ANTISPASMODICS</p> <p>Zanaflex (tizanidine) Flexaril (cyclobenzaprine)</p>
<p style="text-align: center;">AMPHETAMINES (Hold 48 hours prior to exam, resume next day)</p>	<p style="text-align: center;">“ANTI-COAGULANT TYPE DRUGS” (ASA and NSAIDS OK)</p>
<p>Adderol (dextroamphetamine) plus (Amphetamine) Biphetamine (amphetamine) (Amphetamine/CNS Stimulant) Desoxyn (methamphetamine) (Amphetamine/CNS Stimulant) Dexedrine (dextroamphetamine) (Amphetamine/CNS Stimulant) Obetrol (amphetamine) (Amphetamine/CNS Stimulant) Ritalin (methylphenidate) Lisdexamphetamine (Vyvanse)</p>	<p>Coumadin/Warfarin -- Hold 4-5 days prior Plavix/Clopidogrel -- Hold 7 days prior Ticlid/Ticlopidine -- Hold 14 days prior Lovenox/LMWH -- Hold 12 hrs if taking prophylaxis dose (hold > 24 hrs) (also Elmiorn) -- Hold 24 hrs if taking a treatment dose (hold > 24 hrs) Reo-Pro/Abciximab -- Hold 48 hrs prior, given iv only Integrilin/Eptifibatide- Hold 8 hrs prior, given iv only Aggrastat/Tirofiban -- Hold 8hrs prior, given iv only Heparin SC -- If taking >10,000 units, when PTT is normal Cilostazol/Pletal (anti-platelets) – Hold 2 days prior</p> <p><i>MEDICATIONS RESUMED? 24 hrs</i></p>

Medications of concern when Performing Myelograms Dec.2004

Revised Feb. 2005, June 2005, October 2005, November 2005, June 2006, Oct 2006, July 2009 (DKS)